

AD-A107 900

CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY

F/8 11/2

TEMPERATURE DEPENDENCE OF THE HOMOGENEOUS LINEWIDTH OF THE (51D--ETC(U)

NOV 81 J R MORGAN, M A EL-SAYED

N00014-75-C-0602

UNCLASSIFIED

TR-38

NL

1 of 1  
AL  
A107 900



END  
DATE  
FILMED  
1 82  
DTIC

LEVEL II

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0602

Task No. NR 056-498

TECHNICAL REPORT NO. 38

12

AD A107900

TEMPERATURE DEPENDENCE OF THE HOMOGENEOUS LINEWIDTH OF THE  
 $^5D_0 - ^7F_0$  TRANSITION OF  $\text{Eu}^{3+}$  IN AMORPHOUS HOSTS AT HIGH TEMPERATURES

by

Jack R. Morgan and M. A. El-Sayed  
Department of Chemistry  
University of California  
Los Angeles, California 90024

DTIC  
NOV 30 1981  
H

Accepted for Publication

in

Chemical Physics Letters

November 20, 1981

DTIC FILE COPY

Reproduction in whole or in part is permitted for any  
purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

81 11 26 008

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE   |                                      | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM                                  |
|---|--------------------------------------|--|
| 1. REPORT NUMBER<br>Technical Report No. 38   | 2. GOVT ACCESSION NO.<br>AD A107 900 | 3. RECIPIENT'S CATALOG NUMBER  |
| 4. TITLE (and Subtitle)<br>Temperature Dependence of the Homogeneous Line-<br>width of the $^5D_0 - ^7F_0$ Transition of $\text{Eu}^{3+}$ in<br>Amorphous Hosts at High Temperatures  |                                      | 5. TYPE OF REPORT & PERIOD COVERED<br>Interim Technical Report               |
|   |                                      | 6. PERFORMING ORG. REPORT NUMBER   |
| 7. AUTHOR(s)<br>Jack R. Morgan and M. A. El-Sayed   |                                      | 8. CONTRACT OR GRANT NUMBER(s)<br>N00014-75-C-0602                           |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Regents of the University of California<br>University of California, 405 Hilgard Ave.<br>Los Angeles, CA 90024   |                                      | 10. PROGRAM ELEMENT, PROJECT, TASK<br>AREA & WORK UNIT NUMBERS<br>NR-056-498 |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>Office of Naval Research<br>Chemistry Branch<br>Arlington, Virginia 22217  |                                      | 12. REPORT DATE<br>November 20, 1981   |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)<br>Office of Naval Research<br>Branch Office<br>1030 East Green Street<br>Pasadena, CA 91106  |                                      | 13. NUMBER OF PAGES<br>15  |
|   |                                      | 15. SECURITY CLASS. (of this report)<br>unclassified                         |
| 16. DISTRIBUTION STATEMENT (of this Report)<br><br>This document has been approved for public release and sale;<br>distribution of this document is unlimited.  |                                      |  |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  |                                      |  |
| 18. SUPPLEMENTARY NOTES<br><br>Accepted for publication in Chemical Physics Letters   |                                      |  |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br>Glasses<br>Homogenous width<br>Line narrowing<br>Luminescence   |                                      |  |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br>The temperature dependence of the homogeneous linewidth of the $^5D_0 - ^7F_0$<br>transition of $\text{Eu}^{3+}$ in different amorphous solids is determined in the 295°K to<br>800°K temperature range. The nearly quadratic temperature dependence ob-<br>served is discussed in terms of Raman broadening and mechanisms based on two-<br>level systems known to exist in disordered systems. |                                      |  |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TEMPERATURE DEPENDENCE OF THE HOMOGENEOUS LINEWIDTH OF THE  
 $^5D_0 - ^7F_0$  TRANSITION OF  $\text{Eu}^{3+}$  IN AMORPHOUS HOSTS AT HIGH TEMPERATURES


Jack R. Morgan and M. A. El-Sayed

Department of Chemistry

University of California, Los Angeles, California 90024

ABSTRACT

The temperature dependence of the homogeneous linewidth of the  $^5D_0 - ^7F_0$  transition of  $\text{Eu}^{3+}$  in different amorphous solids is determined in the 295 °K to 800 °K temperature range. The nearly quadratic temperature dependence observed is discussed in terms of Raman broadening and mechanisms based on two-level systems known to exist in disordered systems.



## INTRODUCTION

Recently there has been much interest in the origin of the optical homogeneous linewidths of rare-earth ion transitions in amorphous materials. Selzer et al.<sup>1</sup> have observed a nearly quadratic temperature dependence of the homogeneous linewidth of the  $^5D_0 - ^7F_0$  transition of  $\text{Eu}^{3+}$  in a silicate glass from 10 to 100° K. Avouris et al.<sup>2</sup> have reported a similar temperature dependence for the homogeneous linewidth of  $\text{Eu}^{3+}$  in silicate glass in the 200 - 350° K temperature range. Similar results have been obtained for the homogeneous linewidth of  $\text{Pr}^{3+}$  in  $\text{BeF}_2$  and  $\text{GeO}_2$  glasses<sup>3</sup> in the 8 - 300° K range. This behavior of the homogeneous linewidth of rare-earth ions in amorphous hosts cannot be accounted for by the usual one or two phonon broadening mechanisms used to account for the linewidth in crystalline hosts. At temperatures  $T > 100$  K, the homogeneous linewidth in crystalline hosts show a quadratic temperature dependence<sup>1,3</sup>, consistent with a Raman broadening mechanism for  $T > 0.3 T_D$  where  $T_D$  is the Debye temperature. At temperatures below 100 K the Raman contribution to the linewidth drops off as  $T^7$ . For  $T < 100$  K a direct one phonon or Orbach process with a  $T$  or exponential dependence on temperature, respectively, dominate<sup>1,3</sup>. The anomalous low temperature behavior in glass hosts is suggestive of an ion-lattice interaction peculiar to amorphous materials.

Lattice models associated with the disorder inherent to amorphous materials, known as two-level systems (TLS), have been used to interpret many of the low temperature properties of glasses<sup>4,5</sup>. Selzer et al.<sup>1</sup> considered the interaction between pairs of these TLS as a broadening mechanism but predicted a linear temperature dependence. Reinecke<sup>6</sup> has

described a line broadening mechanism involving TLS which also predicts a linear temperature dependence. Lyo and Orbach<sup>7</sup> have described a mechanism involving TLS which predicts a  $T^2$  dependence for temperatures below the Debye temperature. At temperatures above the Debye temperature or cutoff in TLS density of states, whichever is lowest, a linear temperature dependence can be expected<sup>7</sup> for the homogeneous linewidth in glasses.

In order to further understand the homogeneous line broadening mechanism in glasses and to test recent theories we have used fluorescence line narrowing techniques<sup>8</sup> to measure the temperature dependence of the homogeneous linewidth of the  $^5D_0 - ^7F_0$  transition of  $\text{Eu}^{3+}$  in glasses from 295 K to as high as 800 K. Silicate,  $\text{Ca}(\text{PO}_3)_2$ , and  $\text{B}_2\text{O}_3$  glasses with 10, 20, and 30 mole %  $\text{Na}_2\text{O}$  were used. The borate glasses, though similar in composition, are known to undergo significant structural changes<sup>9</sup> with the addition of  $\text{Na}_2\text{O}$ . These structural changes have been shown to affect the spectral properties of  $\text{Eu}^{3+}$ <sup>10,11</sup>. We have found that the homogeneous linewidth follows a nearly quadratic temperature dependence in all of these glasses in the high temperature range. These results, along with those reported in refs. 1-3, show that a smooth quadratic temperature dependence is effective from 10 - 800° K.

#### EXPERIMENTAL

Measurements of the homogeneous linewidth were made on five glass samples. Sample compositions were (mole %): 75  $\text{SiO}_2$ , 15  $\text{Na}_2\text{O}$ , 5  $\text{ZnO}$ , and 5  $\text{BaO}$  for the silicate glass and 100  $\text{Ca}(\text{PO}_3)_2$  for the calcium metaphosphate glasses with 0.25 and 1 mole %  $\text{Eu}_2\text{O}_3$  respectively. The borate sample compositions were (mole %): 90  $\text{B}_2\text{O}_3$ , 10  $\text{Na}_2\text{O}$ ; 80  $\text{Na}_2\text{O}$ , 20  $\text{Na}_2\text{O}$ ;

and 70 B<sub>2</sub>O<sub>3</sub>, 30 Na<sub>2</sub>O with 0.5 Eu<sub>2</sub>O<sub>3</sub> for each of the three samples. Details of the glass preparation are given in ref. 10.

Samples were mounted in an oven for measurements made above room temperature that maintained constant temperature to  $\pm 3^\circ$  K. An Iron-Constantan thermocouple affixed to the sample was used for temperature measurement. For measurements made below room temperature the samples were held in liquid N<sub>2</sub> vapor warmed by a feedback controlled heating core that stabilized the temperature to better than  $\pm 1^\circ$  K.

A Quanta-Ray DCR-1 Nd:YAG pumped PDL-1 pulsed dye laser with a spectral width of  $\sim 0.3 \text{ cm}^{-1}$  and a 6 ns pulse width was used as the excitation source. Excitation was made at 5780 Å for all of the samples, corresponding roughly to the maximum of the inhomogeneous profile of the  $^5\text{D}_0 - ^7\text{F}_0$  transition of Eu<sup>3+</sup> in these glasses. The emission was dispersed with a 1 M Jarrell-Ash scanning monochromator. Resonant detection required the use of a gated phototube<sup>12</sup> to reject scattered excitation light. Signal averaging was accomplished with gated photon counting with a 3 ms window delayed 50  $\mu\text{sec}$  after the laser pulse. In order to assess the contribution of spectral diffusion or other time dependent effects, the detection gate width was decreased from 3 ms to 50  $\mu\text{sec}$  at various temperatures. No change in the observed lineshape was detected on this time scale.

The observed lineshapes for the silicate glass are shown in Fig. 1 for three different temperatures. At room temperature and below the lineshapes are well fit to a Lorentzian. Well above room temperature deviations from a Lorentzian fit were observed on both wings of the line. On the low energy side, overlap with the  $^5\text{D}_0 - ^7\text{F}_1$  fluorescence from resonantly pumped  $^7\text{F}_0 - ^5\text{D}_0$  ions was observed. On the high energy side the intensity falls off more slowly than a Lorentzian. This may be attributed to the  $^5\text{D}_0 - ^7\text{F}_0$  fluorescence of ions pumped from the thermally

populated  ${}^7F_1$  levels via the  ${}^7F_1 - {}^5D_0$  transition. A discussion of this effect is given in ref. 13.

Deconvolution of the instrumental width from the observed width was done using the table of Hegarty et al.<sup>14</sup> to yield an effective measured width ( $\Delta\nu_m$ ). The homogeneous width ( $\Delta\nu_{hom}$ ) was calculated from the measured width using the approximation<sup>15</sup>

$$\Delta\nu_{hom} = \frac{1}{2} \Delta\nu_m$$

for  $\Delta\nu_m < 2/3 \Delta\nu_{inh}$ , where  $\Delta\nu_{inh}$  is the inhomogeneous width of the  ${}^5D_0 - {}^7F_0$  transition which is on the order of 60 - 80  $\text{cm}^{-1}$  fwhm for the glasses used in this study. For  $\Delta\nu_m > 2/3 \Delta\nu_{inh}$ , the tables of ref. 14 were used.

## RESULTS AND DISCUSSION

The results of our measurements of the temperature dependence of the homogeneous linewidth are shown in Fig. 2. The homogeneous linewidth of the  ${}^5D_0 - {}^7F_0$  transition at 5780 Å is found to exhibit a  $T^\alpha$  temperature dependence with  $\alpha$  equal to 2.3, 1.9, 1.9, 2.2, and 2.2 for the silicate, calcium metaphosphate, and borate glasses with 10, 20, and 30 mole %  $\text{Na}_2\text{O}$  respectively; where each of these exponents is good to  $\pm 0.3$ .

The nearly  $T^2$  dependence of the homogeneous linewidth over the measured temperature range is consistent with a two phonon broadening mechanism as has been observed in crystalline hosts<sup>1,3</sup> for  $T > 0.3 T_D$  and conflicts with the high temperature linear dependence predicted by Lyo and Orbach.<sup>7+</sup> In glass hosts, the smooth continuation of the  $T^2$  dependence of the homogeneous linewidth down to 10 K<sup>1,3</sup>, inconsistent with a Raman broadening mechanism based on the Debye model, makes the relative contribution of the Raman mechanism in the high temperature range less clear.

---

<sup>+</sup>These authors arrive at the  $T^2$  dependence by assuming that the density of states of the TLSs as a function of the zero-point energy difference of the two wells is constant on the temperature scale of the experiment. For the temperature range used this doesn't seem reasonable.



Furthermore, the smooth nearly  $T^2$  dependence of the homogeneous linewidth in glasses from 10 K to as high as 800 K, with the same intercept, might suggest that the same mechanism is effective over the entire temperature range.

NMR studies<sup>16,17</sup> of the spin lattice relaxation rate ( $T_1^{-1}$ ) of  $^{11}\text{B}$  and  $^{23}\text{Na}$  in sodium borate and silicate glasses over the 1.2 to 300 K temperature range have shown that the ordinary two phonon Raman contribution was negligible over the entire range investigated. Interpretation of these results<sup>16,17</sup> was also made in terms of processes involving TLS. Recent measurements of the homogeneous linewidths of the  $^5\text{D}_0 - ^7\text{F}_0$  transition of  $\text{Eu}^{3+}$  in sodium borate glasses at room temperature<sup>10</sup> have shown that the homogeneous linewidth increases linearly with the fraction of tetrahedrally coordinated boron atoms. This work also showed a 3.5 power dependence on the velocity of sound,  $v_s$ , for the sodium borate glasses. This is quite different from the  $v_s^{-10}$  dependence expected from the ordinary Raman process<sup>18</sup>. The changes in ion-lattice coupling and type of effective TLS<sup>10</sup> with the structural changes associated with the addition of sodium oxide to borate glass may mask the actual velocity of sound dependence. This might suggest that the homogeneous linewidth in glasses is only weakly dependent on the velocity of sound. Pelligrino et al.<sup>19</sup> have observed a  $\Delta\nu_{\text{hom}} \propto v_s^{-2.6}$  dependence for the homogeneous linewidth of  $\text{Pr}^{3+}$  in a variety of host glasses at room temperature with deviations observed for borate glasses. These results either indicate that the ordinary Raman broadening mechanism is ineffective in determining optical homogeneous linewidths or else the assumptions implied in deriving the dependence on the speed of sound do not hold for glasses.

At the moment, it is difficult to distinguish between two possibilities:

- 1) the TLS model<sup>6,7</sup> is operative in the whole range of temperature but that

theoretical high temperature predictions need further examination or else, 2) the Raman process takes over smoothly<sup>†</sup> (from the TLS mechanism) as the temperature increases without obvious breaks in the  $\Delta\nu_{\text{hom}}$  vs T dependence. It is obvious that more theoretical examination of this problem is needed.

#### ACKNOWLEDGEMENTS

The authors wish to thank Professor R. Orbach for many stimulating discussions and Dr. E. Chock for providing the samples. The financial assistance of the Office of Naval Research is gratefully acknowledged.

---

<sup>†</sup>In the range of  $T \leq 40$  K, R. P. Stout (Ph.D. Dissertation, Iowa State U. 1981) found that a combination of phonon assisted tunnelling and Raman process fit experimental data in reference 3.


## REFERENCES


1. P. M. Selzer, D. L. Huber, D. S. Hamilton, W. M. Yen and M. J. Weber, Phys. Rev. Lett. 36, 813 (1976); P. M. Selzer, D. L. Huber, D. S. Hamilton, W. M. Yen and M. J. Weber, AIP Conference Proceedings No. 31, Structure and Excitation of Amorphous Solids, Williamsburg (1976).
2. P. Avouris, A. Campion and M. A. El-Sayed, J. Chem. Phys. 67, 3397 (1977).
3. J. Hegarty and W. M. Yen, Phys. Rev. Lett. 43, 1126 (1979).
4. P. W. Anderson, B. I. Halperin and C. M. Varma, Phil. Mag. 25, 1 (1972).
5. W. A. Phillips, J. Low Tem. Phys. 7, 351 (1972).
6. T. L. Reinecke, Solid State Commun. 32, 1103 (1979).
7. S. K. Lyo and R. Orbach, Phys. Rev. B 22, 4223 (1980).
8. A. Szabo, Phys. Rev. Lett. 27, 351 (1971).
9. P. J. Bray and T. G. O'Keefe, Phys. Chem. Glasses 4, 37 (1963).
10. J. R. Morgan, E. P. Chock, W. D. Hopewell, M. A. El-Sayed and R. Orbach, J. Phys. Chem. 85, 747 (1981).
11. M. J. Weber, J. Hegarty and D. H. Blackburn, in "Borate Glasses: Structural Properties and Applications," Plenum Press, New York (1978) p. 215.
12. W. Hopewell, thesis, 1980, University of California, Los Angeles.
13. E. Takushi and T. Kushida, J. Lumin. 18/19, 661 (1979).
14. J. Hegarty, R. T. Brondage and W. M. Yen, Appl. Opt. 19, 1889 (1980).
15. T. Kushida and E. Takushi, Phys. Rev. B 12, 824 (1975).
16. J. Szeftel and H. Alloul, Phys. Rev. Lett. 34, 657 (1975).
17. J. Szeftel and H. Alloul, J. Non-Cryst. Solids 29, 253 (1978).
18. B. DiBartolo, Optical Interactions in Solids (John Wiley and Sons, Inc., New York) Ch. 15.
19. J. M. Pelligrino, W. M. Yen and M. J. Weber, J. Appl. Phys. 51, 6332 (1980).

### FIGURE CAPTIONS

Fig. 1. The observed (solid lines) and the Lorentzian fit (solid circles) to the homogeneous lineshapes of the  $^5D_0 - ^7F_0$  transition of  $\text{Eu}^{3+}$  in a silicate glass for resonant emission at  $17300 \text{ cm}^{-1}$  at different temperatures. No change in the observed lineshape was detected upon decreasing the detection gate width from 3 ms to 50  $\mu\text{s}$  thus ruling out spectral diffusion on this time scale.

Fig. 2. Temperature dependence of the homogeneous linewidth ( $\Delta\nu_{\text{hom}}$ ) of the  $^5D_0 - ^7F_0$  emission of  $\text{Eu}^{3+}$  for several glass host observed with resonant excitation at  $17300 \text{ cm}^{-1}$ .  $\circ$  - silicate;  $\Delta$  -  $\text{Ca}(\text{PO}_3)_2$ ;  $\square$  - 70 mol %  $\text{B}_2\text{O}_3$ , 30 mol %  $\text{Na}_2\text{O}$ ;  $\diamond$  - 90 mol %  $\text{B}_2\text{O}_3$ , 10 mol %  $\text{Na}_2\text{O}$ .





TECHNICAL REPORT DISTRIBUTION LIST, GEN

|   | <u>No.</u><br><u>Copies</u> |   | <u>No.</u><br><u>Copies</u> |
|---|-----------------------------|---|-----------------------------|
| Office of Naval Research<br>Attn: Code 472<br>800 North Quincy Street<br>Arlington, Virginia 22217                                    | 2                           | U.S. Army Research Office<br>Attn: CRD-AA-IP<br>P.O. Box 1211<br>Research Triangle Park, N.C. 27709                                 | 1                           |
| ONR Western Regional Office<br>Attn: Dr. R. J. Marcus<br>1030 East Green Street<br>Pasadena, California 91106                         | 1                           | Naval Ocean Systems Center<br>Attn: Mr. Joe McCartney<br>San Diego, California 92152  | 1                           |
| ONR Eastern Regional Office<br>Attn: Dr. L. H. Peebles<br>Building 114, Section D<br>666 Summer Street<br>Boston, Massachusetts 02210 | 1                           | Naval Weapons Center<br>Attn: Dr. A. B. Amster,<br>Chemistry Division<br>China Lake, California 93555                               | 1                           |
| Director, Naval Research Laboratory<br>Attn: Code 6100<br>Washington, D.C. 20390  | 1                           | Naval Civil Engineering Laboratory<br>Attn: Dr. R. W. Drisko<br>Port Hueneme, California 93401                                      | 1                           |
| The Assistant Secretary<br>of the Navy (RE&S)<br>Department of the Navy<br>Room 4E736, Pentagon<br>Washington, D.C. 20350             | 1                           | Department of Physics & Chemistry<br>Naval Postgraduate School<br>Monterey, California 93940  | 1                           |
| Commander, Naval Air Systems Command<br>Attn: Code 310C (H. Rosenwasser)<br>Department of the Navy<br>Washington, D.C. 20360          | 1                           | Scientific Advisor<br>Commandant of the Marine Corps<br>(Code RD-1)<br>Washington, D.C. 20380                                       | 1                           |
| Defense Technical Information Center<br>Building 5, Cameron Station<br>Alexandria, Virginia 22314                                     | 12                          | Naval Ship Research and Development<br>Center<br>Attn: Dr. G. Bosmajian, Applied<br>Chemistry Division<br>Annapolis, Maryland 21401 | 1                           |
| Dr. Fred Saalfeld<br>Chemistry Division, Code 6100<br>Naval Research Laboratory<br>Washington, D.C. 20375                             | 1                           | Naval Ocean Systems Center<br>Attn: Dr. S. Yamamoto, Marine<br>Sciences Division<br>San Diego, California 91232                     | 1                           |
|   |                             | Mr. John Boyle<br>Materials Branch<br>Naval Ship Engineering Center<br>Philadelphia, Pennsylvania 19112                             | 1                           |

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.  
Copies

Mr. James Kelley  
DTNSRDC Code 2803  
Annapolis, Maryland 21402

1

Mr. A. M. Anzalone  
Administrative Librarian  
PLASTEC/ARRADCOM  
Bldg 3401  
Dover, New Jersey 07801

1



TECHNICAL REPORT DISTRIBUTION LIST, DSI

|   | <u>No.</u><br><u>Copies</u> |  | <u>No.</u><br><u>Copies</u> |
|---|-----------------------------|--|-----------------------------|
| Dr. M. A. El-Sayed<br>Department of Chemistry<br>University of California,<br>Los Angeles<br>Los Angeles, California 90024  | 1                           | Dr. M. Rauhut<br>Chemical Research Division<br>American Cyanamid Company<br>Bound Brook, New Jersey 08805              | 1                           |
| Dr. E. R. Bernstein<br>Department of Chemistry<br>Colorado State University<br>Fort Collins, Colorado 80521                 | 1                           | Dr. J. I. Zink<br>Department of Chemistry<br>University of California,<br>Los Angeles<br>Los Angeles, California 90024 | 1                           |
| Dr. C. A. Heller<br>Naval Weapons Center<br>Code 6059<br>China Lake, California 93555                                       | 1                           | Dr. D. Haarer<br>IBM<br>San Jose Research Center<br>5600 Cottle Road<br>San Jose, California 95143                     | 1                           |
| Dr. J. R. MacDonald<br>Chemistry Division<br>Naval Research Laboratory<br>Code 6110<br>Washington, D.C. 20375               | 1                           | Dr. John Cooper<br>Code 6130<br>Naval Research Laboratory<br>Washington, D.C. 20375                                    | 1                           |
| Dr. G. B. Schuster<br>Chemistry Department<br>University of Illinois<br>Urbana, Illinois 61801                              | 1                           | Dr. William M. Jackson<br>Department of Chemistry<br>Howard University<br>Washington, DC 20059                         | 1                           |
| Dr. A. Adamson<br>Department of Chemistry<br>University of Southern<br>California<br>Los Angeles, California 90007          | 1                           | Dr. George E. Walraffen<br>Department of Chemistry<br>Howard University<br>Washington, DC 20059                        | 1                           |
| Dr. M. S. Wrighton<br>Department of Chemistry<br>Massachusetts Institute of<br>Technology<br>Cambridge, Massachusetts 02139 | 1                           |  |                             |

**DAT**  
**ILM**